

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

APOC1, 27-83aa, Human, His tag, E.coli

产品货号: TP01195

第三版

别名:Apolipoprotein C-I precursor , APOC1B, Apo-CIB, ApoC-IB, Apolipoprotein C-I, Apolipoprotein C1

描述:APOC1 is a member of the apolipoprotein C family. This gene is expressed primarily in the liver, and it is activated when monocytes differentiate into macrophages. A pseudogene of this gene is located 4 kb downstream in the same orientation, on the same chromosome. It is mapped to chromosome 19, where it resides within an apolipoprotein gene cluster. Alternatively spliced transcript variants have been found for this gene, but the biological validity of some variants has not been determined. APOC1 has a length of 57 amino acids normally found in plasma and responsible for the activation of esterified lecithin cholesterol with an important role in the exchange of esterified cholesterol between lipoproteins and in removal of cholesterol from tissues. Its main function is inhibition of CETP, probably by altering the electric charge of HDL molecules. Recombinant human APOC1 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

配方:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 10% glycerol.

分子量:9 kDa (80aa) confirmed by MALDI-TOF

序列:

MGSSHHHHHHSSGLVPRGSHMGSTPDVSSALDKLKEFGNTLEDKARELISRIKQSELSAKMREWFSETFQKVKEKLKIDS

纯度:> 95% by HPLC

浓度:0.25 mg/ml (determined by Bradford assay)

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.