

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

BAG1, 1-230aa, Human, E.coli

产品货号: TP01290

第三版

别名: BCL2-associated athanogene 1, HAP, RAP46.

描述: BCL2-associated athanogene, also known as BAG1, is a member of the Bcl-2 family of proteins that play varied stimulatory or inhibitory roles in the regulation of programmed cell death. This functions in conjunction with Bcl-2 to inhibit apoptosis.

配方: Liquid. In 20mM Tris-HCl buffer (pH7.5) containing 100mM NaCl, 10% glycerol

分子量: 25.9 kDa (230aa), confirmed by MALDI-TOF

序列:

MNRSQEVTRDEESTRSEEVTR EEMAAAGLTVTVTHSNEKHDLHVTSQQGSSEPVVQDLAQVVEEVIGVPQSFQKLIFKGK
SLKEMETPLSALGIQDGCRVMLIGKKNSPQEEVELKHLKLEKSVEKIADQLEELNKELTGIQQGFLPKDLQAEALCKLDRR
VKATIEQFMKILEEIDTLILPENFKDSRLKRKGLVKKVQAFLAECTVEQNICQETERLQSTNFALAE

纯度: > 95% by HPLC

浓度: 1 mg/ml (determined by Bradford assay)

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.