

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

BCL10, 1-233aa, Human, His tag, E.coli

产品货号: TP01310

第三版

别名: B-cell lymphoma/leukemia 10, c-E10, CARMEN, CIPER, CLAP, mE10.

描述: BCL10, also known as B-cell lymphoma/leukemia 10, contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF-kappaB. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF-kappaB signaling.

配方: Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol, 0.05M NaCl, 1mM DTT

分子量: 28.8 kDa (257aa) confirmed by MALDI-TOF

序列:

MGSSHHHHHHSSGLVPRGSHMGSHMEPTAPSLTEEDLTVKDALENLRVYLCEKIIAERHFDHLRAKKILSREDTEEISC
RTSSRKRAKLLDYLQENPKGLDTLVESIRREKTQNFLIQKITDEVLKLRLNIKLEHLKGLKCSSCEPFPDGATNNLSRSNSDES
NFSEKLRASVVMYHPEGESSTPPFFSTNSSLNLPVLEVGRTEENTIFSSSTLPRPGDPGAPPLPPDLQLEEEGTCANSSEMFLPLR
SRTVSRQ

纯度: > 95% by HPLC

浓度: 0.5 mg/ml (determined by Bradford assay)

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.