

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

BCL7C, 1-217aa, Human, His tag, E.coli

产品货号: TP01316

第三版

别名: B-cell CLL/lymphoma 7 protein family member C,

描述: BCL7C is identified by the similarity of its product to the N-terminal region of BCL7A protein. The BCL7A protein is encoded by the gene known to be directly involved in a three-way gene translocation in a Burkitt lymphoma cell line. The function of this gene has not yet been determined. Recombinant human BCL7C protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques

配方: Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 20% glycerol, 1mM DTT

分子量: 25.9kDa (240aa) confirmed by MALDI-TOF (Molecular size on SDS-PAGE will appear higher)

序列:

MGSSHHHHHSSGLVPRGSHMGSMAGRTVRAETRSRAKDDIKKVMATIEKVRWWEKRWVTVGDTSLRIFKWVPPVDP  
QEEERRRAGGGAERSRGRERRGRGASPRGGGPLILLDLNDENSNQSFHSEGLQKGTEPSPGGTPQPSRPVSPAGPPEGVPE  
EAQPPRLGQERDPGGITAGSTDEPPMLTKEEPVPELLEAEAPEAYPVFEPVPPVPEAAQGDTEDESEGAPPLKRICPNAPDP

纯度: > 95% by HPLC

浓度: 0.25 mg/ml (determined by Bradford assay)

内毒素: <1.0 EU per 1 ug of protein (determined by LAL method)

存储: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.