

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

EIF3I, 1-325aa, Human, His tag, E.coli

产品货号: TP01995

第三版

别名:eukaryotic translation initiation factor 3 subunit I, C3orf68

描述:EIF3I is a component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2: GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Recombinant human EIF3I protein, fused to His-tag at N-terminus, was expressed in E.coli.

配方:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol 0.4M Urea

分子量:38.9kDa (348aa)

序列:

MGSSHHHHHSSGLVPRGSHMGSMKPILLQGHHERSITQIKYNREGDLLFTVAKDPIVNVWYSVNGERLGTYMGHTGAV
WCVDADWDTKHVLTGSADNSCRLWDCETGKQLALLKTNSAVRTCGFDGFGNIIMFSTDKQMGYQCFVSFFDLRDP
SQI
DNNEPYMKIPCNDKITSVWGPLGECIIAGHESGELNQYSAKSGEVLVNVKEHSRQINDIQLSRDMTMFV
TASKDNTAK
LFDSTTLEHQKTFRTERPVNSAALSPNYDHVVLGGGQEAMDVTTTSTRIGKFEARFFHLAFEEEEFGRV
KGFHFGPINSVAFH
PDGKSYSSGGEDGYVRIHYFDPQYFEFEFEA

纯度:> 95% by HPLC

浓度:1.0 mg/ml (determined by Bradford assay)

内毒素:<1.0 EU per 1 ug of protein (determined by LAL method)

存储:+4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.