

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

APOH, 20-345aa, Human, His tag, E.coli

Cat.NO.: TP01196

3th Edition

Synonyms: Beta-2-glycoprotein 1, Apolipoprotein H, B2G1, B2GP1, BG

Description: Apolipoprotein H, also known as APOH, binds to various kinds of negatively charged substances such as heparin, phospholipids, and dextran sulfate. This protein may prevent activation of the intrinsic blood coagulation cascade by binding to phospholipids on the surface of damaged cells.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 2M urea, 10% glycerol

Molecular Weight: 38.6 kDa (349aa)

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSGRTCPKPDDLPFSTVVPLKTFYEPGEEITYSCKPGYVSRGGMRKFICPLTGL WPINTLKCTPRVCPFAGILENGAVRYTTFEYPNTISFSCNTGFYLNGADSAKCTEEGKWSPELPVCAPIICPPPSIPT FATLRVYKPSAGNNSLYRDTAVFECLPQHAMFGNDTITCTTHGNWTKLPECREVKCPFPSRPDNGFVNYPAKPTL YYKDKATFGCHDGYSLDGPEEIECTKLGNWSAMPSCKASCKVPVKKATVVYQGERVKIQEKFKNGMLHGDKVSFF CKNKEKKCSYTEDAQCIDGTIEVPKCFKEHSSLAFWKTDASDVKPC

Purity:> 95% by HPLC

Concentration: 0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1