

CD40LG, 113-261aa, Human, His tag, E.coli

Cat.NO.: TP01516

3th Edition

Synonyms:CD40 ligand, CD154, CD40L, gp39; hCD40L, HIGM1, IGM, IMD3, T-BAM, TNFSF5, TRAP

Description:CD40 ligand, also known as CD40LG, is a member of the TNF superfamily and is expressed on activated T cells. CD40LG is expressed as a soluble cytokine as well as a homotrimeric type II transmembrane protein. It has been reported to be important for B cell costimulation following binding of its receptor, CD40.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 40% glycerol, 0.2M NaCl

Molecular Weight:18.3 kDa (169aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMQKGDQNPQIAAHVISEASSKTTSVLQWAEKGYTMSNNLVTLENGKQLTVKRQ
GLYYIYAQVTFCNREASSQAPFIASLCLKSPGRFERILLRAANTHSSAKPCGQQSIHLGGVFELQPGASVFNVT
PSQVSHGTGFTSFGLLKL

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.