

CD8B, 22-170aa, Human, His tag, E.coli

Cat.NO.: TP01540

3th Edition

Synonyms:T-cell surface glycoprotein CD8 beta chain isoform 5 precursor, CD8B1, LEU2, LY3, LYT3, P37

Description:CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T-cell receptor on the T lymphocyte recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 beta chain isoforms. Recombinant human CD8B protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M UREA, 10% glycerol

Molecular Weight:19.2kDa (172aa)

Sequences:

MGSSHHHHHSSGLVPRGSHMGSLQQTPAYIKVQTNKMVMLSCEAKISLSNMRIYWLRQRQAPSSDSHHEFLAL
WDSAKGTIHGEEVEQEKI AVFRDASRFILNLT SVKPEDSGIYFCMIVGSPELTFGKGTQLSVVDLPTTAQPTKKSTL
KKRVCRLPRPETQKGPLCSP

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.