

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

CDCA8, 19-280aa, Human, His tag, E.coli

Cat.NO.: TP01549

3th Edition

Synonyms: Cell division cycle-associated protein 8, Borealin, Dasra-B, PESCRG3

**Description:**CDCA8, also known as Borealin, is a component of the chromosomal passenger complex (CPC). The CPC complex has essential functions at the centromere in ensuring correct chromosome alignment and segregation and is required for chromatin-induced microtubule stabilization and spindle assembly.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT,50% glycerol, 1mM EDTA, 0.2M NaCl

Molecular Weight: 31.6 kDa (283aa) confirmed by MALDI-TOF

## Sequences:

MGSSHHHHHHSSGLVPRGSHMRKLASFLKDFDREVEIRIKQIESDRQNLLKEVDNLYNIEILRLPKALREMNWLDYF ALGGNKQALEEAATADLDITEINKLTAEAIQTPLKSAKTRKVIQVDEMIVEEEEEEENERKNLQTARVKRCPPSKKRT QSIQGKGKGKRSSRANTVTPAVGRLEVSMVKPTPGLTPRFDSRVFKTPGLRTPAAGERIYNISGNGSPLADSKEIFL TVPVGGGESLRLLASDLQRHSIAQLDPEALGNIKKLSNRLAQICSSIRTHK

Purity:> 95% by HPLC

Concentration: 0.25 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1