

CDK2AP2, 1-126aa, Human, His tag, E.coli

Cat.NO.: TP01559

3th Edition

Synonyms: cyclin-dependent kinase 2 associated protein 2, DOC-1R, p14

Description: CDK2AP2, also known as DOC-1R and p14, belongs to the CDK2AP family. CDK2-associated protein is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. CDK2AP2 is a regulator for self-renewal of mouse embryonic stem cells (mESCs) under permissive conditions, and cell survival during differentiation of the mESCs into terminally differentiated cell types. CDK2AP2 may be involved in the regulation of self-renewal of stem cells during early embryogenesis. Recombinant human CDK2AP2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form: Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.2M NaCl, 50% glycerol, 2mM DTT

Molecular Weight: 15.5kDa (149aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGMSYKPIAPAPSSTPGSSTPGPGTPVPTGSVPSPSGSVPGAGAPFRPLFNDF
GPPSMGYVQAMKPPGAQGSQSTYTDLLSVIEEMGKEIRPTYAGSKSAMERLKRGIHARALVRECLAETERNART

Purity: > 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level: <1.0 EU per 1 ug of protein (determined by LAL method)

Storage: Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.