

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

COX5A, 42-150aa, Human, His tag, E.coli

Cat.NO.: TP01694

3th Edition

Synonyms:cytochrome c oxidase subunit Va, COX, COX-VA, VA

**Description:**COX5A, also known as COX, COX-VA and VA, belongs to the cytochrome c oxidase subunit 5A family. Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. COX5A is the heme A-containing chain of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport. Recombinant human COX5A protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 10% glycerol, 2mM DTT

Molecular Weight: 14.9kDa (132aa), confirmed by MALDI-TOF

## Sequences:

MGSSHHHHHHSSGLVPRGSHMGSSHGSQETDEEFDARWVTYFNKPDIDAWELRKGINTLVTYDMVPEPKIIDAAL RACRRLNDFASTVRILEVVKDKAGPHKEIYPYVIQELRPTLNELGISTPEELGLDKV

Purity:> 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1