

CSDC2, 1-153aa, Human His tag, E.coli**Cat.NO.: TP01736**

3th Edition

Synonyms: Cold shock domain-containing protein C2, dJ347H13.2, PIPPIN

Description: CSDC2, as known as cold shock domain-containing protein C2, is RNA-binding factor which binds specifically to the very 3'-UTR ends of both histone H1 and H3.3 mRNAs, encompassing the polyadenylation signal. The cold shock domain containing proteins (CSDPs) are one group of the evolutionarily conserved nucleic acid-binding proteins widely distributed in bacteria, plants, animals, and involved in various cellular processes, including adaptation to low temperature, cellular growth, nutrient stress and stationary phase. It may play a central role in the negative regulation of histone variant synthesis in the developing brain. Recombinant human CSDC2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography.

Form: Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol 0.1M NaCl

Molecular Weight: 19.2 kDa(176aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSMSTSESTSPPVVPLHSPKSPVWPTFPFHHREGSRVWERGGVPPRDLPSPLP
TKRTRTYSATARASAGPVFKGVCKQFSRSQGHGFITPENGSEDIFVHVSDIEGEYVPVEGDEVITYKMCPIPPKNQK
FQAVEVVLTLQAPHTPHETWSGQVVGS

Purity: > 95% by HPLC

Concentration: 0.5 mg/ml (determined by Bradford assay)

Endotoxin Level: <1.0 EU per 1 ug of protein (determined by LAL method)

Storage: Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.