

DTD1, 1-209aa, Human, His tag, E.coli

Cat.NO.: TP01928

3th Edition

Synonyms:D-tyrosyl-tRNA deacylase 1, bA379J5.3, bA555E18.1, C20orf88, DUEB, HARS2, pqn-68

Description:DTD1 hydrolyzes D-tyrosyl-tRNA(Tyr) into D-tyrosine and free tRNA(Tyr). It could be a defense mechanism against a harmful effect of D-tyrosine. DTD1 may be a risk factor for aspirin-intolerant asthma (AIA) by catalyzing the hydrolysis of D-tryptophan and interacting with the tyrosyl-tRNA synthetase (tyrRS) enzyme, which promotes a pro-inflammatory phenotype.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 10% glycerol, 1mM DTT

Molecular Weight:25.9kDa (232aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSMKAVVQRVTRASVTVGGEQISAIGRGICVLLGISLEDTQKELEHMVRKILNLR
VFEDESGKHWSKSVMDKQYEILCVSQFTLQCVLKGKPDFHLAMPTEQAEGFYNSFLEQLRKTYRPELIKDGKFGA
YMQVHIQNDGPVTIELESPAPGTATSDPKQLSKLEKQQQRKEKTRAKGPSESSKERNTPRKEDRSASSGAEGDVS
SEREP

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.