

Recombinant Human HDAC8 / HDACL1 Protein (GST tag)

Cat.NO.: TP07230

3th Edition

Synonyms:CDA07;CDLS5;HD8;HDACL1;MRXS6;RPD3;WTS

Description:Histone deacetylase 8, also known as HDAC8 and HDACL1, is a nucleus and cytoplasm protein which belongs to the histone deacetylase family and HD type 1 subfamily. Histone deacetylases (HDACs) are a growing family of enzymes implicated in transcriptional regulation by affecting the acetylation state of core histones in the nucleus of cells. HDAC8 / HDACL1 is weakly expressed in most tissues. It expressed at higher level in heart, brain, kidney and pancreas and also in liver, lung, placenta, prostate and kidney. HDAC8 / HDACL1 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. HDAC8 / HDACL1 may play a role in smooth muscle cell contractility. HDAC8 / HDACL1 may be a potential drug target for neuroblastoma differentiation therapy using selective inhibitors, avoiding unspecific side effects.

Form:PBS

Molecular Weight:68 kDa

Sequences:Met 1-Val 377

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.