

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human LAIR1 Protein (His Tag)

Cat.NO.: TP07251

3th Edition

Synonyms:CD305;LAIR-1

Description:Leukocyte associated Ig-like receptor-1 (LAIR1) is a surface molecule expressed on human mononuclear leukocytes that functions as an inhibitory receptor on human NK cells. In addition to NK cells, LAIR1 is expressed on T cells, B cells, macrophages, and dendritic cells. It is predicted to mediate inhibitory functions based on the presence of immunoreceptor tyrosine-based inhibitory motifs (ITIMs) in its cytoplasmic domain. Crosslinking of LAIR1 on human T cell clones results in inhibition of cytotoxicity only in T cell clones that lack CD28 and are able to spontaneously lyse certain targets in vitro. Moreover, the cytolytic activity of freshly isolated T cells, which is thought to be mainly due to "effector" T cells, can be inhibited by anti-LAIR1 mAb. Thus, LAIR1 functions as an inhibitory receptor not only on NK cells, but also on human T cells. This indicates that LAIR1 provides a mechanism of regulation of effector T cells and may play a role in the inhibition of unwanted bystander responses mediated by Ag-specific T cells.

Form:PBS

Molecular Weight: 17.2 kDa

Sequences: Met 1-Tyr165

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1 / 1