
Recombinant Mouse ENTPD5 Protein (His tag)

Cat.NO.: TP07584

3th Edition

Synonyms:AI196558;AI987697;Cd39l4;ER-UDPase;mNTPase;NTPDase-5;NTPDase5;Pcph

Description:Ectonucleoside triphosphate diphosphohydrolase 5 (ENTPD5), also known as CD39 antigen-like 4, ER-UDPase, Guanosine-diphosphatase ENTPD5, Nucleoside diphosphatase Uridine-diphosphatase ENTPD5. This hydrolase is expressed in response to phosphoinositide 3-kinase (PI3K) signaling. Activation of PI3K results in FOXO phosphorylation by AKT1 and loss of ENTPD5 transcriptional repression. It is Up-regulated in PTEN-deficient cells. Uridine diphosphatase (UDPase) that promotes protein N-glycosylation and ATP level regulation.ENTPD5 promotes protein N-glycosylation and folding in the endoplasmic reticulum, as well as elevated ATP consumption in the cytosol via an ATP hydrolysis cycle. Together with CMPK1 and AK1, ENTPD5 constitutes an ATP hydrolysis cycle that converts ATP to AMP and results in a compensatory increase in aerobic glycolysis. ENTPD5 also hydrolyzes GDP and IDP but not any other nucleoside di-, mono- or triphosphates, nor thiamine pyrophosphate. This enzyme Plays a key role in the AKT1-PTEN signaling pathway by promoting glycolysis in proliferating cells in response to phosphoinositide 3-kinase (PI3K) signaling.

Form:PBS

Molecular Weight:46.6 kDa

Sequences:Thr19-Ser427

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.