

---

**Recombinant Human TCN2 Protein (His tag)**

**Cat.NO.: TP07591**

3th Edition

---

**Synonyms:**D22S676;D22S750;II;TC;TC-2;TC2;TCII

**Description:**Transcobalamin II, also known as TCN2 and TC II, is a plasma protein that binds cobalamin (Cbl; vitamin B12) as it is absorbed in the terminal ileum and distributes to tissues. The circulating transcobalamin II-cobalamin complex binds to receptors on the plasma membrane of tissue cells and is then internalized by receptor-mediated endocytosis. Transcobalamin II is a non-glycosylated secretory protein of molecular mass 43 kDa. Its plasma membrane receptor (TC II-R) is a heavily glycosylated protein with a monomeric molecular mass of 62 kDa. Human TCN2 gene is composed of nine exons and eight introns spanning approximately 20 kb with multiple potential transcription start sites. A number of genetic abnormalities are characterized either by a failure to express TCN2 or by synthesis of an abnormal protein. The TCN2 deficiency results in cellular cobalamin deficiency, an early onset of megaloblastic anaemia, and neurological abnormalities.

**Form:**PBS

**Molecular Weight:**46.7 kDa

**Sequences:**Met 1-Trp 427

**Purity:**> 95% by HPLC

**Concentration:**

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.