
Recombinant Mouse JAM2 / CD322 / VE-JAM Protein (His tag)

Cat.NO.: TP07593

3th Edition

Synonyms:1110002N23Rik;2410030G21Rik;2410167M24Rik;JAM-2;JAM-B;Jcam2;VE-JAM

Description:Junctional adhesion molecule B (JAM-B), also known as Junctional adhesion molecule 2 (JAM2), Vascular endothelial junction-associated molecule (VE-JAM), and CD322, is a single-pass type I membrane protein which belongs to the immunoglobulin superfamily. It is prominently expressed on high endothelial venules. expression to be restricted to the high endothelial venule of tonsil and lymph nodes. The localization to the endothelium of arterioles in and around inflammatory and tumor foci. JAM-B can function as an adhesive ligand for the T cell line J45 and can interact with GM-CSF/IL-4-derived peripheral blood dendritic cells, circulating CD56(+) NK cells, circulating CD56(+)CD3(+) NK/T cells, and circulating CD56(+)CD3(+)CD8(+) cytolytic T cells. JAM-2 is expressed on high endothelial venules (HEVs) in human tonsil and on a subset of human leukocytes, suggesting that JAM-2 plays a central role in the regulation of transendothelial migration. It binds to very late activation antigen (VLA)-4, a leucocyte integrin that contributes to rolling and firm adhesion of lymphocytes to endothelial cells through binding to vascular cell adhesion molecule (VCAM)-1. JAM-B appears to contribute to leucocyte extravasation by facilitating not only transmigration but also rolling and adhesion. JAM-B acts as an adhesive ligand for interacting with a variety of immune cell types and may play a role in lymphocyte homing to secondary lymphoid organs.

Form:PBS

Molecular Weight:24.7 kDa

Sequences:Met 1-Asn 236

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.