

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human ACY1 / Aminoacylase-1 Protein (His tag)

Cat.NO.: TP07613

3th Edition

Synonyms: ACY-1; ACY1D; HEL-S-5

Description:Aminoacylase 1 (ACY1), a metalloenzyme that removes amide-linked ACY1 groups from amino acids and may play a role in regulating responses to oxidative stress. Both the C-terminal fragment found in the two-hybrid screen and full-length ACY1 co-immunoprecipitate with SphK1. Though both C-terminal and full-length proteins slightly reduce SphK1 activity measured in vitro, the C-terminal fragment inhibits while full-length ACY1 potentiates the effects of SphK1 on proliferation and apoptosis. It suggested that ACY1 physically interacts with SphK1 and may influence its physiological functions. As a homodimeric zinc-binding enzyme, Aminoacylase 1 catalyzes the hydrolysis of N alpha-acylated amino acids. Deficiency of Aminoacylase 1 due to mutations in the Aminoacylase 1 (ACY1) gene follows an autosomal-recessive trait of inheritance and is characterized by accumulation of N-acetyl amino acids in the urine.

Form:PBS

Molecular Weight: 47.3 kDa

Sequences:Met 1-Ser 408

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1