

**Recombinant Human Cathepsin V / Cathepsin L2 / Preproprotein Protein (His tag)****Cat.NO.: TP07630**

3th Edition

**Synonyms:**CATL2;CTSL2;CTSU;CTSV;MGC125957

**Description:**Cathepsin V (CTSV), also known as Cathepsin L2, CTSL2, and CATL2, is a member of the peptidase C1 family. It is predominantly expressed in the thymus and testis. Cathepsin V is also expressed in corneal epithelium, and to a lesser extent in conjunctival epithelium and skin. It is a lysosomal cysteine proteinase that may play an important role in corneal physiology. It has about 75% protein sequence identity to murine cathepsin L. The fold of this enzyme is similar to the fold adopted by other members of the papain superfamily of cysteine proteases. Cathepsin V has been recently described as highly homologous to Cathepsin L and exclusively expressed in human thymus and testis. Cathepsin V is the dominant cysteine protease in cortical human thymic epithelial cells, while Cathepsin L and Cathepsin S seem to be restricted to dendritic and macrophage-like cells. Active Cathepsin V in thymic lysosomal preparations was demonstrated by active-site labeling. Recombinant Cathepsin V was capable of converting I $\kappa$ B into CLIP efficiently, suggesting that it is the protease that controls the generation of I $\kappa$ B-CLIP complexes in the human thymus. Cathepsin V is the third elastolytic cysteine protease which exhibits the most potent elastase activity yet described among human proteases and that it is present in atherosclerotic plaque specimens. Cathepsin L2 may play a specialized role in the thymus and testis. Expression analysis of cathepsin L2 in human tumors revealed a widespread expression in colorectal and breast carcinomas but not in normal colon or mammary gland or in peritumoral tissues. Cathepsin L2 was also expressed by colorectal and breast cancer cell lines as well as by some tumors of diverse origin, including ovarian and renal carcinomas.

**Form:**PBS**Molecular Weight:**37.1 kDa**Sequences:**Met 1-Val 334**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1  $\mu$ g of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.