

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human LAG3 Protein (Fc Tag)

Cat.NO.: TP07698

3th Edition

Synonyms:CD223;LAG-3

Description:LAG3, also known as CD223 and Lymphocyte activation gene 3, belongs to immunoglobulin (Ig) superfamily. LAG3 contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. It is selectively expressed in activated T and NK cells. LAG3 has a negative regulatory function in T cells. It also acts as as a new marker of T cell induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signalling, leading to increased antigen-specific T-cell responses in vivo.Immune CheckpointTargets Co-inhibitory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy

Form:PBS

Molecular Weight:72.3 kDa

Sequences: Met 1-Leu450

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1