
Recombinant Human MFG-E8 / lactadherin / MFGE8 Protein (His tag)**Cat.NO.: TP07725**

3th Edition

Synonyms:BA46;EDIL1;HMFG;hP47;HsT19888;MFG-E8;MFGM;OAcGD3S;SED1;SPAG10

Description:MFG-E8, also known as lactadherin and MFGE8, contains 1 EGF-like domain and 2 F5/8 type C domains. It also contains a phosphatidylserine (PS) binding domain, as well as an Arginine-Glycine-Aspartic acid motif, which enables the binding to integrins. It binds PS, which is exposed on the surface of apoptotic cells. MFG-E8 is expressed in mammary epithelial cell surfaces and aortic media. Overexpression of MFG-E8 can be found in several carcinomas. MFG-E8 has an opsonization of the apoptotic cells and binding to integrins on the surface of phagocytic cells. It also mediates the engulfment of the dead cell. MFG-E8 plays an important role in the maintenance of intestinal epithelial homeostasis and the promotion of mucosal healing. It promotes VEGF-dependent neovascularization and contributes to phagocytic removal of apoptotic cells in many tissues. It also binds to phosphatidylserine-enriched cell surfaces in a receptor-independent manner.

Form:PBS**Molecular Weight:**42 kDa**Sequences:**Met 1-Cys 387**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.