

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Cystatin A / CSTA Protein (His tag)

Cat.NO.: TP07747

3th Edition

Synonyms: AREI; STF1; STFA

Description:Cystatin-A, also known as Cystatin-AS, Stefin-A and CSTA, is a cytoplasm protein which belongs to the cystatin family. Cystatin-A / CSTA is a cysteine proteinase inhibitor with a molecular mass of 11 kDa, and is located mainly in the keratohyaline granules of the stratum granulosum and the cornified envelope of the stratum corneum in the epidermis. The cystatins are a family of cysteine protease inhibitors with homology to chicken cystatin. Cystatins are physiological inhibitors of cysteine proteinases which are widely distributed in human tissues and fluids. Cystatins typically comprise about 115 amino acids, are largely acidic, contain four conserved cysteine residues known to form two disulfide bonds. Cystatins may be glycosylated and / or phosphorylated, with similarity to fetuins, kininogens, stefins, histidine-rich glycoproteins and cystatin-related proteins. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired inhibitory activity. Cystatins mainly inhibit peptidases belonging to peptidase families C1 (papain family) and C13 (legumain family).

Form:PBS

Molecular Weight: 12.4 kDa

Sequences: Ile 2-Phe 98

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1 / 1