

---

**Recombinant Human VSIG4 Protein (Fc tag)****Cat.NO.: TP07872**

3th Edition

**Synonyms:**CR1g;Z391g

**Description:**VSIG4 (V-set and immunoglobulin domain containing 4), also known as complement receptor of the immunoglobulin superfamily (CR1g) and Z391g, is a type I transmembrane glycoprotein. It is a B7 family-related protein and an Ig superfamily member. In contrast to the B7 family members which contain two IgG domains, VSIG4 contains one complete V-type I g domain and a truncated C-type I g domain. VSIG4 is exclusively expressed on tissue resident macrophages and binds to multimers of C3b and iC3b that are covalently attached to particle surfaces. No VSIG4 expression appears to be present in T and B cells. VSIG4 functions as a negative regulator of T cell activation, and may be involved in the maintenance of peripheral T cell tolerance, and is also identified as a potent suppressor of established inflammation. Mouse VSIG4 is synthesized as a 280 amino acid precursor that contains a signal sequence, an V-type I g domain (aa 36-115), one potential N-linked glycosylation site, and a single transmembrane domain. The V-type I g domain of mouse VSIG4 shares 86% and 80% aa sequence identity with the V-type I g domains of rat and human VSIG4, respectively.

**Form:**PBS**Molecular Weight:**56.2 kDa**Sequences:**Met 1-Pro283**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.