

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human ULBP1 / N2DL1 Protein (His tag)

Cat.NO.: TP07894

3th Edition

Synonyms: RAET11

Description: UL16-binding proteins (ULBP) or retinoic acid early transcripts-1 (RAET1) are ligands to the activating receptor, NKG2D. Ten members of the human ULBP/RAET1 gene family have been identified to encode for potentially functional proteins, and have tissue-specific expressions. ULBP1, also known as RAET1I and NKG2DL1, together with at least ULBP 2 and 3, are well-known ligands for NKG2D, and activate multiple signaling pathways in primary NK cells, resulting in the production of cytokines and chemokines. ULBP1 is expressed in T-cells, B-cells, erythroleukemia cell lines and in a wide range of tissues including heart, brain, lung, liver and bone marrow, as well as some tumor cells. As an unconventional member of the MHC class I family, ULBP1 function in immune responses, especially in cancer and infectious diseases. Unlike other ULBP members, ULBP1 is able to interact with soluble CMV glycoprotein UL16 in CMV infected cells. The interaction with UL16 blocked the interaction with the NKG2D receptor, and thus might escape the immune surveillance. Furthermore, UL16 also causes ULBP1 to be retained in the ER and cis-Golgi apparatus so that it does not reach the cell surface. The ULBP1 regulation may have implications for development of new therapeutic strategies against cancer cells.

Form:PBS

Molecular Weight:23.8 kDa

Sequences: Met 1-Gly 216

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1