
Recombinant Human IL20RA Protein (His tag)**Cat.NO.: TP07921**

3th Edition

Synonyms:CRF2-8;IL-20R-alpha;IL-20R1;IL-20RA

Description: Interleukin 20 receptor, alpha subunit (IL20RA/IL-20RA) also known as IL-20 receptor subunit alpha, Cytokine receptor class-II member 8, interleukin-20 receptor I, and interleukin-20 receptor subunit alpha, is a subunit for the interleukin-20 receptor. IL20RA/IL-20RA belongs to the type II cytokine receptor family. This cytokine is a receptor for interleukin 20 (IL20), a cytokine that may be involved in epidermal function. The receptor of IL20 is a heterodimeric receptor complex consisting of this protein and interleukin 20 receptor beta (IL20B). IL20RA forms heterodimer with IL20RB, and the complex serves as a receptor for IL19, IL20 and IL24. All three are capable of signaling through IL-20RA/IL-20RB complex. The ligand binding to receptor B creating a high-affinity binding site for the receptor A which is recruited to complete the complex. In addition, IL20RA also forms a heterodimer with the unique and specific receptor IL10RB and functions as the receptor for IL26. IL20RA is widely expressed with highest levels in skin, testis and brain. The expression of both IL20RA and IL20RB is found to be upregulated in psoriatic skin lesions on keratinocytes.

Form:PBS**Molecular Weight:**26.8 kDa**Sequences:**Met 1-Lys 250**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.