

Recombinant Human CCDC47 Protein (His tag)

Cat.NO.: TP08055

3th Edition

Synonyms:GK001;MSTP041

Description:CCDC47 gene is expressed at high level. The gene contains 16 distinct gt-ag introns. Transcription produces 9 different mRNAs, 6 alternatively spliced variants and 3 unspliced forms. There are 3 probable alternative promoters, 3 non overlapping alternative last exons and 8 validated alternative polyadenylation sites. The mRNAs appear to differ by truncation of the 5' end, truncation of the 3' end, presence or absence of a cassette exon, overlapping exons with different boundaries. Functionally, CCDC47 gene has been proposed to participate in processes such as calcium ion homeostasis, embryo development, ER overload response and post-embryonic development. CCDC47 are expected to have molecular function (calcium ion binding) and to localize in various compartments (membrane, endoplasmic reticulum, integral to membrane, microsome).

Form:PBS

Molecular Weight:14.7 kDa

Sequences:Met 1-Ser135

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.