

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Contactin 2 / CNTN2 Protein (His tag)

Cat.NO.: TP08061

3th Edition

Synonyms: AXT; FAME5; TAG-1; TAX; TAX1

Description: Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and NB-3. Since their identification in the late 1980s, Contactin-1 and Contactin-2 have been studied extensively. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. In contrast, the function of the other four members of this subgroup remained unknown until recently. Contactin-2, also known as CNTN2, is a glycosylphosphatidylinositol (GPI)-anchored neuronal membrane protein that functions as a cell adhesion molecule. The human, rat, and chicken Contactin-2 are alternatively known as TAX1 (transiently-expressed axonal glycoprotein), TAG1 (transient axonal glycoprotein), and axonin-1, respectively. Human Contactin-2 shares approximately 91% and 75% amino acid sequence identity with rat and chicken Contactin-2, respectively. Contactin-2 is expressed by a subset of neuronal populations in the developing central nervous system (CNS) and peripheral nervous system (PNS). Contactin-2 is also expressed by oligodendrocytes and Schwann cells, which are myelinating glial cells of the CNS and PNS, respectively. Contactin-2 may play a role in the formation of axon connections in the developing nervous system. Contactin-2 is also involved in glial tumorigenesis and may provide a potential target for therapeutic intervention. During embryonic development, Contactin-2 interacts either in a homophilic, or heterophilic fashion with various transmembrane proteins.

Form: PBS

Molecular Weight: 109 kDa

Sequences: Met 1-Asn 1012

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1