

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Mouse c-MET / HGFR Protein (His tag)

Cat.NO.: TP08342

3th Edition

Synonyms:Al838057;c-Met;HGF;HGFR;Par4

Description:Hepatocyte growth factor receptor (HGFR), also known as c-Met or mesenchymal-epithelial transition factor (MET), is a receptor tyrosine kinase (RTK) that has been shown to be overexpressed and/or mutated in a variety of malignancies. HGFR protein is produced as a single-chain precursor, and HGF is the only known ligand. Normal HGF/HGFR signaling is essential for embryonic development, tissue repair or wound healing, whereas aberrantly active HGFR has been strongly implicated in tumorigenesis, particularly in the development of invasive and metastatic phenotypes. HGFR protein is a multifaceted regulator of growth, motility, and invasion, and is normally expressed by cells of epithelial origin. Preclinical studies suggest that targeting aberrant HGFR signaling could be an attractive therapy in cancer.

Form:PBS

Molecular Weight: 102 kDa

Sequences:Met 1-Asn 929

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1