

**Recombinant Human B7-1 Protein****Cat.NO.: TP08396**

3th Edition

**Synonyms:**B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB7

**Description:**The B-lymphocyte activation antigen B7-1 (referred to as B7), also known as CD80, is a member of cell surface immunoglobulin superfamily and is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. As costimulatory ligands, B7-1 which exists predominantly as dimer and the related protein B7-2, interact with the costimulatory receptors CD28 and cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expressed on T cells, and thus constitute one of the dominant pathways that regulate T cell activation and tolerance, cytokine production, and the generation of CTL. The B7/CD28/CTLA4 pathway has the ability to both positively and negatively regulate immune responses. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.

**Form:**PBS**Molecular Weight:**25.5 kDa**Sequences:**Met 1-Asn 242**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.