

Recombinant Human B7-DC / PD-L2 / CD273 Protein (His tag)**Cat.NO.: TP08583**

3th Edition

Synonyms:B7-DC;B7DC;bA574F11.2;Btdc;CD273;PD-L2;PDCD1L2;PDL2

Description:Programmed death ligand 2 (PD-L2), also referred to as B7-DC and CD273, is a member of the B7 family of proteins including B7-1, B7-2, B7-H2, B7-H1 (PD-L1), and B7-H3. PD-L2 is a type I membrane protein and structurally consists of an extracellular region containing one V-like and one C-like Ig domain, a transmembrane region, and a short cytoplasmic domain. PD-L2 is expressed on antigen presenting cells, placental endothelium and medullary thymic epithelial cells, and can be induced by LPS in B cells, INF- γ in monocytes, or LPS plus IFN- γ in dendritic cells. The CD28 and B7 protein families are critical regulators of immune responses. PD-L2 and PD-L1 are two ligands for PD-1, member of the CD28/CTLA4 family expressed on activated lymphoid cells, and thus provide signals for regulating T cell activation and immune tolerance. The interaction of B7-DC/PD-1 exhibited a 2-6-fold higher affinity compared with the interaction of B7-H1/PD-1.

Immune Checkpoint
Detection: Antibodies **Immune Checkpoint Detection:** ELISA Antibodies **Immune Checkpoint Detection:** FCM Antibodies **Immune Checkpoint Detection:** WB Antibodies
Immune Checkpoint Proteins
Immune Checkpoint Targets **Co-inhibitory Immune Checkpoint Targets** **Immunotherapy** **Cancer Immunotherapy** **Targeted Therapy**

Form:PBS**Molecular Weight:**24 kDa**Sequences:**Met 1-Pro 219**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.