

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## Recombinant Human IDO1 / IDO Protein (His tag)

Cat.NO.: TP08608

3th Edition

Synonyms:IDO;IDO-1;INDO;Indoleamine 2,3â€'dioxygenase

Description:Indoleamine 2,3-dioxygenase-1, also known as Indoleamine-pyrrole 2,3-dioxygenase, IDO1 and IDO, is a member of the indoleamine 2,3-dioxygenase family. IDO1 / IDO and tryptophan 2,3-dioxygenase (TDO) are tryptophan-degrading enzymes that catalyze the first step in tryptophan catabolism via the kynurenine pathway. TDO is widely distributed in both eukaryotes and bacteria. In contrast, IDO has been found only in mammals and yeast. In 2007, a third enzyme, indoleamine 2,3-dioxygenase-2 (IDO2), was discovered. IDO2 is found not only in mammals but also in lower vertebrates. IDO1 / IDO is an immunosuppressive molecule inducible in various cells. IDO1 / IDO catalyzes the cleavage of the pyrrol ring of tryptophan and incorporates both atoms of a molecule of oxygen. It mediates oxidative cleavage of tryptophan, an amino acid essential for cell proliferation and survival. IDO1 / IDO inhibition is proposed to have therapeutic potential in immunodeficiency-associated abnormalities, including cancer. The IDO pathway is activated in multiple tumor types. Selective inhibition of IDO1 may represent an attractive cancer therapeutic strategy via up-regulation of cellular immunity. IDO1 / IDO is an enzyme that suppresses adaptive T-cell immunity by catabolizing tryptophan from the cellular microenvironment. Inhibition of IDO pathway might enhance the efficacy of immunotherapeutic strategies for cancer.Immune CheckpointImmune Checkpoint Detection: ELISA AntibodiesCo-inhibitory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy

Form:PBS

Molecular Weight:46 kDa

Sequences: Ala 2-Gly 403

Purity:> 95% by HPLC

**Concentration:** 

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1