

Recombinant Human CD20 / MS4A1 Protein (TrxA Tag)**Cat.NO.: TP08632**

3th Edition

Synonyms:B1;Bp35;CD20;CVID5;LEU-16;MS4A1;MS4A2;S7

Description:CD20 (membrane-spanning 4-domains, subfamily A, member 1), also known as MS4A1, is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. CD20 / MS4A1 is expressed on all stages of B cell development except the first and last. CD20 / MS4A1 is present from pre-pre B cells through memory cells, but not on either pro-B cells or plasma cells. It is a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. CD20 / MS4A1 may be involved in the regulation of B-cell activation and proliferation. Defects in CD20 / MS4A1 are the cause of immunodeficiency common variable type 5 (CVID5). CVID5 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B-cells is usually in the normal range, but can be low. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

Form:PBS**Molecular Weight:**23.9 kDa**Sequences:**Ile141-Ser188**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.