
本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Smad2L/3L (T220/T179 Phosphorylated) (1A1) Mouse IgG MoAb

单克隆抗体

产品货号: MA01589

第三版

免疫原: Synthetic peptide of phosphorylated Smad2L/3L (T220/T179).

亚型: IgG1

克隆: 1A1

描述: Phosphorylation of signal transduction molecules, Smad2 and Smad3, can be an important information for understanding of various biological functions of transforming growth factor (TGF)- β . TGF- β type I receptor and cyclin-dependent kinases phosphorylate both Smad2 and Smad3 at the C-terminals and at linker (middle) regions of them respectively (ref. 1). TGF-signaling is mediated by Smad phosphoisoforms phosphorylated at both the C-terminal and linker regions (ref. 2, 3). This monoclonal antibody against for Smad2L/3L (T220/T179) recognizes the part of phosphorylated threonin (Thr220 of Smad2 and Thr179 of Smad3) located in linker region of Smad2/3 specifically. It can be used for western blotting, immunoprecipitation and immunohistochemistry. It is applicable for enzyme biochemical analysis of TGF- β signaling and makes it possible to visualize the intermolecular reaction of phosphorylated Smad signaling in human tissues and monitor them in real-time. Thus, analysis of TGF- β signaling with this antibody is expected to be widely applied to cancer serearch (ref. 4) and fibrosis research (ref. 5), and to contribute to understanding the wide variety of life phenomina mediated by phosphoisoforms of Smad2/3.

配方: Lyophilized product in PBS containing 1 % BSA and 0.05 % NaN₃

浓度: Please see datasheet.

保存: +4 ° C 保存 (1-2 周). 长期保存在 -20 ° C 或者 -70 ° C. 避免反复冻融.

应用: ELISA, WB, ICC/IF