

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Anti-Human ERBB2 Monoclonal Antibody

单克隆抗体

产品货号: MA01788

第三版

免疫原:Synthetic Peptide

亚型:IgG

克隆:2F1

描述:Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization. In the nucleus is involved in transcriptional regulation. Associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. Implicated in transcriptional activation of CDKN1A; the function involves STAT3 and SRC. Involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.

配方:

浓度:1mg/mL

+4°C保存(1-2周).长期保存在-20°C或者-70°C.避免反复冻融.

应用:ELISA, WB, ICC/IF

1/1