

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human CX3CL1 Monoclonal Antibody

单克隆抗体

产品货号: MA02213

第三版

免疫原:Fusion protein of CX3CL1

亚型:IgG1

克隆:357

**描述:**This gene encodes a large cytokine protein of 373 amino acids, it contains multiple domains and is the only known member of the CX3C chemokine family. It is also commonly known under the names fractalkine (in humans) and neurotactin (in mice). The polypeptide structure of CX3CL1 differs from the typical structure of other chemokines. CX3CL1 is produced as a long protein (with 373-amino acid in humans) with an extended mucin-like stalk and a chemokine domain on top. The mucin-like stalk permits it to bind to the surface of certain cells. However a soluble (90 kD) version of this chemokine has also been observed. Soluble CX3CL1 potently chemoattracts T cells and monocytes, while the cell-bound chemokine promotes strong adhesion of leukocytes to activated endothelial cells, where it is primarily expressed. CX3CL1 elicits its adhesive and migratory functions by interacting with the chemokine receptor CX3CR1. Its gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

**配方:**

浓度:None

保存: +4 ° C 保存 (1-2 周). 长期保存在-20 ° C或者-70 ° C. 避免反复冻融.

应用:ELISA, WB, ICC/IF