

本公司提供的电子版本说明书仅供参考,实验请以收到的纸质手册为准。

Anti-Human/Mouse/Rat Acetyl-Histone H2A (Lys15) Polyclonal Antibody

多克隆抗体

产品货号: PA01460

第三版

描述:HIST1H2AG (Histone Cluster 1 H2A Family Member G) is a Protein Coding gene. Among its related pathways are Activated PKN1 stimulates transcription of AR (androgen receptor) regulated genes KLK2 and KLK3 and Deubiquitination. GO annotations related to this gene include sequence-specific DNA binding and enzyme binding. An important paralog of this gene is HIST1H2AM. HIST2H2AA3 (Histone Cluster 2 H2A Family Member A3) is a Protein Coding gene. Diseases associated with HIST2H2AA3 include Retinitis Pigmentosa 20. Among its related pathways are Activated PKN1 stimulates transcription of AR (androgen receptor) regulated genes KLK2 and KLK3 and Meiosis. GO annotations related to this gene include sequence-specific DNA binding and protein heterodimerization activity. An important paralog of this gene is HIST2H2AA4. HIST3H2A (Histone Cluster 3 H2A) is a Protein Coding gene. Among its related pathways are Activated PKN1 stimulates transcription of AR (androgen receptor) regulated genes KLK2 and KLK3 and Deubiquitination. GO annotations related to this gene include sequence-specific DNA binding and protein heterodimerization activity. An important paralog of this gene is HIST2H2AA4. HIST3H2A (Histone Cluster 3 H2A) is a Protein Coding gene. Among its related pathways are Activated PKN1 stimulates transcription of AR (androgen receptor) regulated genes KLK2 and KLK3 and Deubiquitination. GO annotations related to this gene include sequence-specific DNA binding and protein heterodimerization activity. An important paralog of this gene is HIST1H2AC.

抗原:Synthetic Peptide

配方:

如何使用:加1ml超纯水重溶

稳定性: -20°C保存条件下,冻干粉,保质期为五年;液体,保质期为两年。

稀释液:PBS (pH7.4) , 1% BSA

应用:WB1~5µg/ml.

特异性: