

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human/Mouse/Rat Phospho-CHEK1 (Ser317) Polyclonal Antibody

多克隆抗体

产品货号: PA01527

第三版

描述: Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by proteolysis of cyclins. Chk1 and Chk2 are involved in these processes as regulators of Cdks. Chk1 and Chk2 both function as essential components in the G2 DNA damage checkpoint by phosphorylating Cdc25C in response to DNA damage. Phosphorylation inhibits Cdc25C activity, thereby blocking mitosis. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. It has also been shown that Chk1 can phosphorylate Wee 1 in vitro, providing evidence that the hyperphosphorylated form of Wee 1, seen in cells delayed by Chk1 overexpression, is due to phosphorylation by Chk1.

抗原: Synthesized peptide derived from human Chk1 around the phosphorylation site of Ser317

配方:

如何使用: 加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液: PBS (pH7.4) ， 1% BSA

应用: WB 1 ~ 5 μg/ml.

特异性: Expressed ubiquitously with the most abundant expression in thymus, testis, small intestine and colon.