

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human/Mouse/Rat Phospho-EGFR (Tyr1197) Polyclonal Antibody

多克隆抗体

产品货号: PA01548

第三版

描述: Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses. Known ligands include EGF, TGFA/TGF-alpha, amphiregulin, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin. Isoform 2 may act as an antagonist of EGF action.

抗原: Synthesized peptide derived from human EGFR around the phosphorylation site of Tyr1197

配方:

如何使用: 加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液: PBS (pH7.4) ， 1% BSA

应用: WB 1 ~ 5 μ g/ml.

特异性: Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.