

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human BAG1 Polyclonal Antibody

多克隆抗体

产品货号: PA02322

第三版

**描述:**The oncogene BCL2 is a membrane protein that blocks a step in a pathway leading to apoptosis or programmed cell death. The protein encoded by this gene binds to BCL2 and is referred to as BCL2-associated athanogene. It enhances the anti-apoptotic effects of BCL2 and represents a link between growth factor receptors and anti-apoptotic mechanisms. Multiple protein isoforms are encoded by this mRNA through the use of a non-AUG (CUG) initiation codon, and three alternative downstream AUG initiation codons. A related pseudogene has been defined on chromosome X.

**抗原:**Synthesized peptide derived from the Internal region of human Bag-1.

**配方:**

**如何使用:**加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:**PBS (pH7.4) ， 1% BSA

**应用:**WB 1 ~ 5  $\mu$ g/ml.

**特异性:**Isoform 4 is the most abundantly expressed isoform. It is ubiquitously expressed throughout most tissues, except the liver, colon, breast and uterine myometrium. Isoform 1 is expressed in the ovary and testis. Isoform 4 is expressed in several types of tumor cell lines, and at consistently high levels in leukemia and lymphoma cell lines. Isoform 1 is expressed in the prostate, breast and leukemia cell lines. Isoform 3 is the least abundant isoform in tumor cell lines (at protein level).