

本公司提供的电子版说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human /Mouse CST4 Polyclonal Antibody

多克隆抗体

产品货号: PA02841

第三版

描述:The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a type 2 salivary cysteine peptidase inhibitor.

抗原:Recombinant protein of human CST4

配方:

如何使用:加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液:PBS (pH7.4) ， 1% BSA

应用:WB 1 ~ 5 μ g/ml.

特异性:Expressed in submandibular and sublingual saliva but not in parotid saliva (at protein level). Expressed in saliva, tears, urine and seminal fluid.