

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human/Mouse/Rat Phospho-MERTK/TYRO3 (Tyr753/Tyr685) Polyclonal Antibody

多克隆抗体

产品货号: PA04470

第三版

**描述:**MERTK is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in this gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP).MERTK (MER Proto-Oncogene, Tyrosine Kinase) is a Protein Coding gene. Diseases associated with MERTK include Retinitis Pigmentosa 38 and Mertk-Related Retinitis Pigmentosa. Among its related pathways are GPCR Pathway and Nanog in Mammalian ESC Pluripotency. GO annotations related to this gene include transferase activity, transferring phosphorus-containing groups and protein tyrosine kinase activity. An important paralog of this gene is AXL. TYRO3 (TYRO3 Protein Tyrosine Kinase) is a Protein Coding gene. Diseases associated with TYRO3 include External Ear Carcinoma and Lymphocytic Choriomeningitis. Among its related pathways are GPCR Pathway and Nanog in Mammalian ESC Pluripotency. GO annotations related to this gene include transferase activity, transferring phosphorus-containing groups and protein tyrosine kinase activity. An important paralog of this gene is AXL.

**抗原:**Synthetic peptide from human protein

**配方:**

**如何使用:**加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:**PBS (pH7.4) ， 1% BSA

**应用:**WB 1 ~ 5  $\mu$ g/ml.

**特异性:**MERTK is Not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines. Highly expressed in testis, ovary, prostate, lung, and kidney, with lower expression in spleen, small intestine, colon, and liver. TYRO3 is Abundant in the brain and lower levels in other tissues.