

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human CLEC4A Polyclonal Antibody

多克隆抗体

产品货号: PA04517

第三版

**描述:** This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 transmembrane protein may play a role in inflammatory and immune response. Multiple transcript variants encoding distinct isoforms have been identified for this gene. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer gene complex region.

**抗原:** Synthesized peptide derived from CLECSF6

**配方:**

**如何使用:** 加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:** PBS (pH7.4) , 1% BSA

**应用:** WB 1 ~ 5 μ g/ml.

**特异性:** Expressed in dendritic cells, myeloid cells, B-cells and HL-60 cells (at protein level). TNF alpha, IL-1 alpha, and LPS, down-regulated expression at the surface of neutrophils (at protein level). Expressed preferentially in hematopoietic tissues. Expressed in peripheral blood leukocytes, neutrophils, moderate quantities in spleen, lymph node, and bone marrow, and at very low levels in thymus. Expressed in Ag-presenting cells (DC, monocytes, macrophages and B-cells), as well as on granulocytes. Expression was decreased in DC by signals inducing its maturation (e.g. CD40 ligand, LPS, and TNF alpha).