

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human IL17RC Polyclonal Antibody

多克隆抗体

产品货号: PA04606

第三版

描述: This gene encodes a single-pass type I membrane protein that shares similarity with the interleukin-17 receptor (IL-17RA). Unlike IL-17RA, which is predominantly expressed in hemopoietic cells, and binds with high affinity to only IL-17A, this protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. The proinflammatory cytokines, IL-17A and IL-17F, have been implicated in the progression of inflammatory and autoimmune diseases. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, and it has been proposed that soluble, secreted proteins lacking transmembrane and intracellular domains may function as extracellular antagonists to cytokine signaling.

抗原: Synthesized peptide derived from the C-terminal region of human IL-17RC

配方:

如何使用: 加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液: PBS (pH7.4) ， 1% BSA

应用: WB 1 ~ 5 μg/ml.

特异性: Expressed in prostate, skeletal muscle, kidney and placenta (at protein level) (PubMed:11706037). Expressed in brain, cartilage, colon, heart, intestine, kidney, liver, lung, muscle, placenta, and prostate (PubMed:11706037). Also detected in thyroid, trachea and adrenal gland (PubMed:17911633). Low expression in thymus and leukocytes (PubMed:11706037).