

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human/Mouse/Rat MNAT1 Polyclonal Antibody

多克隆抗体

产品货号: PA04844

第三版

描述:Stabilizes the cyclin H-CDK7 complex to form a functional CDK-activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIH basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminus domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II.

抗原:Synthesized peptide derived from the Internal region of human Mat1

配方:

如何使用:加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液:PBS (pH7.4) ， 1% BSA

应用:WB 1 ~ 5 μ g/ml.

特异性:Highest levels in colon and testis. Moderate levels are present thymus, prostate, ovary, and small intestine. The lowest levels are found in spleen and leukocytes.