

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human ZnT-8 Polyclonal Antibody

多克隆抗体

产品货号: PA06186

第三版

描述:The protein encoded by this gene is a zinc efflux transporter involved in the accumulation of zinc in intracellular vesicles. This gene is expressed at a high level only in the pancreas, particularly in islets of Langerhans. The encoded protein colocalizes with insulin in the secretory pathway granules of the insulin-secreting INS-1 cells. Allelic variants of this gene exist that confer susceptibility to diabetes mellitus, noninsulin-dependent (NIDDM). Several transcript variants encoding different isoforms have been found for this gene.SLC30A8 (Solute Carrier Family 30 Member 8) is a Protein Coding gene. Diseases associated with SLC30A8 include Diabetes Mellitus, Noninsulin-Dependent and Prediabetes Syndrome. Among its related pathways are Transport of glucose and other sugars, bile salts and organic acids, metal ions and amine compounds and Metal ion SLC transporters. GO annotations related to this gene include protein homodimerization activity and zinc ion transmembrane transporter activity. An important paralog of this gene is SLC30A2.

抗原:Synthesized peptide derived from the Internal region of human ZnT-8

配方:

如何使用:加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液:PBS (pH7.4) ， 1% BSA

应用:WB 1 ~ 5 μ g/ml.

特异性:In the endocrine pancreas, expressed in insulin-producing beta cells. Expressed at relatively high levels in subcutaneous fat tissue from lean persons; much lower levels in visceral fat, whether from lean or obese individuals, and in subcutaneous fat tissue from obese individuals. Expressed in peripheral blood mononuclear cells, including T-cells and B-cells, with great variation among individuals ranging from negative to strongly positive.