

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human CD32-A Polyclonal Antibody

多克隆抗体

产品货号: PA06391

第三版

**描述:**CD32 (also designated Fc gamma RII) is a low affinity receptor for the Fc fragment of aggregated IgG (1,2). CD32 is responsible for the clearance of immunocomplexes by macrophages and also plays an important role in the regulation of antibody production by B cells (1-4). IgG can noncooperatively bind either one or two highly glycosylated CD32 molecules, and this binding delivers a negative signal for B cells (1,2,5). CD32 exists as several isoforms that are produced by alternative splicing of three distinct genes, A, B, and C (2,6). These isoforms are designated FcγRIIA, FcγRIIB1, FcγRIIB3, and FcγRIIC (1,2,6). All isoforms are present on monocytes, placental trophoblasts and endothelial cells (1,6). In addition, the FcγRIIB forms are present on B lymphocytes, and the FcγRIIA and FcγRIIC forms are found on neutrophils (1,6).

**抗原:**Synthesized peptide derived from the Internal region of human CD32-A.

**配方:**

**如何使用:**加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:**PBS (pH7.4) ， 1% BSA

**应用:**WB 1 ~ 5 μg/ml.

**特异性:**Found on monocytes, neutrophils and eosinophil platelets.