

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human C4A/B Polyclonal Antibody

多克隆抗体

产品货号: PA06479

第三版

**描述:**C4A (Complement C4A (Rodgers Blood Group)) is a Protein Coding gene. Diseases associated with C4A include C4a Deficiency and Immunodeficiency Due To A Classical Component Pathway Complement Deficiency. Among its related pathways are Complement Pathway and Innate Immune System. GO annotations related to this gene include endopeptidase inhibitor activity and complement component C1q binding. An important paralog of this gene is C4B. C4B (Complement C4B (Chido Blood Group)) is a Protein Coding gene. Diseases associated with C4B include C4b Deficiency and Immunodeficiency Due To A Classical Component Pathway Complement Deficiency. Among its related pathways are Innate Immune System and Complement Pathway. GO annotations related to this gene include carbohydrate binding and complement binding. An important paralog of this gene is C4A.

**抗原:**Synthesized peptide derived from the Internal region of human C4a/b.

**配方:**

**如何使用:**加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:**PBS (pH7.4) ， 1% BSA

**应用:**WB 1 ~ 5 μ g/ml.

**特异性:**Complement component C4 is expressed at highest levels in the liver, at moderate levels in the adrenal cortex, adrenal medulla, thyroid gland,and the kidney, and at lowest levels in the heart, ovary, small intestine, thymus, pancreas and spleen. The extra-hepatic sites of expression may be important for the local protection and inflammatory response.1 Publication