
本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human EMR2 Polyclonal Antibody

多克隆抗体

产品货号: PA06544

第三版

描述: This gene encodes a member of the class B seven-span transmembrane (TM7) subfamily of G-protein coupled receptors. These proteins are characterized by an extended extracellular region with a variable number of N-terminal epidermal growth factor-like domains coupled to a TM7 domain via a mucin-like spacer domain. The encoded protein is expressed mainly in myeloid cells where it promotes cell-cell adhesion through interaction with chondroitin sulfate chains. This gene is situated in a cluster of related genes on chromosome 19. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. ADGRE2 (Adhesion G Protein-Coupled Receptor E2) is a Protein Coding gene. Diseases associated with ADGRE2 include Vibratory Urticaria and Physical Urticaria. Among its related pathways are Presynaptic function of Kainate receptors and Peptide ligand-binding receptors. GO annotations related to this gene include G-protein coupled receptor activity and transmembrane signaling receptor activity. An important paralog of this gene is ADGRE5.

抗原: Synthesized peptide derived from the Internal region of human EMR2.

配方:

如何使用: 加1ml超纯水重溶

稳定性: -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

稀释液: PBS (pH7.4) ， 1% BSA

应用: WB 1 ~ 5 μ g/ml.

特异性: Expression is restricted to myeloid cells. Highest expression was found in peripheral blood leukocytes, followed by spleen and lymph nodes, with intermediate to low levels in thymus, bone marrow, fetal liver, placenta, and lung, and no expression in heart, brain, skeletal muscle, kidney, or pancreas. Expression is also detected in monocyte/macrophage and Jurkat cell lines but not in other cell lines tested.