

本公司提供的电子版本说明书仅供参考，实验请以收到的纸质手册为准。

Anti-Human MDFIC Polyclonal Antibody

多克隆抗体

产品货号: PA10395

第三版

**描述:** This gene product is a member of a family of proteins characterized by a specific cysteine-rich C-terminal domain, which is involved in transcriptional regulation of viral genome expression. Alternative translation initiation from an upstream non-AUG (GUG), and an in-frame, downstream AUG codon, results in the production of two isoforms, p40 and p32, respectively, which have different subcellular localization; p32 is mainly found in the cytoplasm, whereas p40 is targeted to the nucleolus. Both isoforms have transcriptional regulatory activity that is attributable to the cysteine-rich C-terminal domain. Alternative splicing results in multiple transcript variants.

**抗原:** Synthetic peptide of human MDFIC

**配方:**

**如何使用:** 加1ml超纯水重溶

**稳定性:** -20 ° C保存条件下，冻干粉,保质期为五年；液体，保质期为两年。

**稀释液:** PBS (pH7.4) ， 1% BSA

**应用:** WB 1 ~ 5 μ g/ml.

**特异性:** Expressed in lymphoid organs (spleen, thymus, peripheral blood leukocytes) as well as prostate, uterus and small intestine. 1  
Publication